| 5. | A curve | is | described | by | the | equation |
|-----------|---------|----|-----------|----|-----|----------|
|-----------|---------|----|-----------|----|-----|----------|

$$x^3 - 4y^2 = 12xy.$$

(a) Find the coordinates of the two points on the curve where x = -8.

(3)

(b) Find the gradient of the curve at each of these points.

(6)

| Leave | |
|-------|--|
| blank | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

| 4. A curve has equation $3x^2 - y^2 + xy = 4$. The points <i>P</i> and <i>Q</i> lie of the tangent to the curve is $\frac{8}{3}$ at <i>P</i> and at <i>Q</i> . | on the curve. The gradient |
|--|----------------------------|
| (a) Use implicit differentiation to show that $y - 2x = 0$ at P and | nd at Q. (6) |
| (b) Find the coordinates of P and Q . | (3) |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

| A curve C has the equation $y^2 - 3y = x^3 + 8$. | |
|---|-----|
| (a) Find $\frac{dy}{dx}$ in terms of x and y. | |
| dx | (4) |
| | () |
| (b) Hence find the gradient of C at the point where $y = 3$. | (3) |
| | (3) |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

| Leave |
|-------|
| hlank |

| 4. | The curve C has the equation $ye^{-2x} = 2x + y^2$. | |
|----|--|--|
| | | |

(a) Find $\frac{dy}{dx}$ in terms of x and y.

(5)

The point P on C has coordinates (0, 1).

(b) Find the equation of the normal to C at P, giving your answer in the form ax + by + c = 0, where a, b and c are integers.

(4)

| |
|------|
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |



Leave

3. The curve C has the equation

$$\cos 2x + \cos 3y = 1$$
, $-\frac{\pi}{4} \leqslant x \leqslant \frac{\pi}{4}$, $0 \leqslant y \leqslant \frac{\pi}{6}$

(a) Find $\frac{dy}{dx}$ in terms of x and y.

(3)

The point *P* lies on *C* where $x = \frac{\pi}{6}$.

(b) Find the value of y at P.

(3)

(c) Find the equation of the tangent to C at P, giving your answer in the form $ax + by + c\pi = 0$, where a, b and c are integers.

(3)

| Question 3 continued | blank |
|----------------------|-------|
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |



| Leave | |
|-------|--|
| blank | |

| A curve C has equation $2^x + y^2 = 2xy$ | |
|--|-----|
| Find the exact value of $\frac{dy}{dx}$ at the point on C with coordinates (3, 2). | |
| | (7) |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |